

# HEDIS 2015 - Cervical Cancer Screening (CCS)

The percentage of women 21–64 years of age who were screened for cervical cancer using either of the following criteria: Women age 21–64 who had cervical cytology performed every 3 years OR Women age 30–64 who had cervical cytology/human papillomavirus (HPV) co-testing performed every 5 years.

Improvement Notation	Higher score indicates better quality
Denominator	Women 24–64 years as of the last day of the measurement year.
Numerator	Identify women 24–64 years of age as of the last day of the measurement year who had cervical cytology (Cervical Cytology Value Set) during the measurement year or the two years prior to the measurement year OR women 35–64 years of age as of the last day of of the measurement year who had cervical cytology (Cervical Cytology Value Set) and a human papillomavirus (HPV) test (HPV Tests Value Set) with service dates four or less days apart during the measurement year or the four years prior to the measurement year.
Exclusions	Hysterectomy with no residual cervix (Hysterectomy Value Set) any time during the member’s history through the last day of the measurement year.
Exceptions	None
Note	

Last updated 01/13/2017

## Reference Periods Reference Date 02/12/2017

Base Period	02/12/2016 - 02/12/2017	12 months prior to the reference date
Ever	- 02/12/2017	Any time prior to the reference date
Reporting Period Cervicalcytology	02/12/2014 - 02/12/2017	2 years prior to start date of Base Period
Reporting Period Hpv	02/12/2013 - 02/12/2017	3 years prior to start date of Base Period


Denominator

Between 24 and 64 years of age on end date of **Base Period** — 02/12/2017

AND

Gender is Female

Numerator



Exclusion

☐

One code indicating the following for **Ever** — any time before 02/12/2017

✓ Hysterectomy

See table **XCL1**

Reference Tables

---

Denominator Reference Tables

Numerator Reference Tables


N1	✔ Cervical cytology	
	Observation	
	LOINC	
	10524-7, 18500-9, 19762-4, 19764-0, 19765-7, 19766-5, 19774-9, 33717-0, 47527-7, 47528-5	
	Procedure	
	CPT	HCPCS
	88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88154, 88164, 88165, 88166, 88167, 88174, 88175	G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, P3000, P3001, Q0091

N2	✔ HPV testing	
	Observation	
	LOINC	
	21440-3, 30167-1, 38372-9, 49896-4, 59420-0	
	Procedure	
	CPT	
	87620, 87621, 87622	

Exclusions Reference Tables

XCL1	✔ Hysterectomy	
	Diagnosis	
	ICD-9	
	618.5, 752.43, V88.01, V88.03	
	Procedure	
	CPT	ICD-9
	51925, 56308, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290, 58291, 58292, 58293, 58294, 58548, 58550, 58552, 58553, 58554, 58570, 58571, 58572, 58573, 58951, 58953, 58954, 58956, 59135	68.4, 68.41, 68.49, 68.5, 68.51, 68.59, 68.6, 68.61, 68.69, 68.7, 68.71, 68.79, 68.8

Reference Periods



Different criteria in each measure require that records occur during a precise time period. Note that all dates begin at midnight.

Operators

Operators describe what criteria is needed for a measure. They include:



Both criteria above and below the line must be met



Either criteria above or below the line must be met



The criteria following this are negated — they must not have occurred

Expressions

You will encounter certain expressions that denote a specific relationship between criteria. They include:



Linked Encounter  
Indicates that the criteria listed above must have occurred during the same encounter as the criteria that immediately follows



Indicates that the criteria must have occurred in a certain sequence. The criteria that follows must have occurred after the criteria listed above



Reverse sequence — the criteria that follows must have occurred before the criteria listed above



The first criteria specifies what must be on the patient record. The criteria below is then counted for that patient