

CHP Custom - Rolling Preventive Care and Screening: Screening for Clinical Depression and Follow-Up Plan

Percentage of patients aged 12 years and older screened for clinical depression during the measurement period.

Improvement Notation	Higher score indicates better quality
Denominator	All patients aged 12 years and older before the beginning of the measurement period with at least one eligible encounter during the measurement period
Numerator	Patients screened for clinical depression on the date of the encounter using an age appropriate standardized tool AND if positive, a follow-up plan is documented on the date of the positive screen
Exclusions	Patients that expired or entered hospice during the reporting period. Patients with an active diagnosis for Depression or a diagnosis of Bipolar Disorder
Exceptions	Patient Reason(s): Patient refuses to participate OR Medical Reason(s): Patient is in an urgent or emergent situation where time is of the essence and to delay treatment would jeopardize the patient's health status OR Situations where the patient's functional capacity or motivation to improve may impact the accuracy of results of standardized depression assessment tools. For example: certain court appointed cases or cases of delirium
Note	

Last updated 01/13/2017

Reference Periods

Reference Date 02/12/2017

Base Period	02/12/2016 - 02/12/2017	12 months prior to the reference date
Ever	- 02/12/2017	Any time prior to the reference date
Reporting Period Lookback	02/12/2015 - 02/12/2017	1 year prior to start date of Base Period
Reporting Period Dayprior	02/11/2016 - 02/12/2016	1 day prior to start date of Base Period and 12 months prior to end date

For help reading this document, refer to key on last page.

Denominator

All codes indicating the following for **Base Period** — between 02/12/2016 and 02/12/2017

See table D1



Happening during same encounter as the following

Happening during same encounter as the following

✓ PREV-12 Encounter, Outpatient or Office Visit

See table L1

AND

At least 12 years of age on start date of **Base Period** — 02/12/2016

Numerator

One code indicating the following

depression screening

See table N1

Exclusion

	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017	
OR	✓ Deceased	See table XCL1
	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=20)	
	✓ Deceased	See table XCL2
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=40)	
	✓ Deceased	See table XCL2
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=41)	
	✓ Deceased	See table XCL2
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=42)	
	✓ Deceased	See table XCL2
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017	
	✓ Hospice	See table XCL3
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=50)	
	✓ Hospice	See table XCL2
OR	One code indicating the following for Base Period — between 02/12/2016 and 02/12/2017 (dischargeDisposition=51)	
	✓ Hospice	See table XCL2
OR	One code indicating the following for Reporting Period Dayprior — between 02/11/2016 and 02/12/2016 (sourceDataList=PROBLEM_LIST)	
	✓ Dx of Depression or Bipolar Disorder in the Problem List	See table XCL4
OR	One code indicating the following, on at least 2 different days for Reporting Period Lookback — between 02/12/2015 and 02/12/2017	
	✓ Dx of Depression or Bipolar Disorder on 2 encounters in the prior year	See table XCL4

Exception

One code indicating the following for **Base Period** — between 02/12/2016 and 02/12/2017

✓ Medical or Patient Exception

See table **XCE1**

Reference Tables

Linked Encounter Reference Tables

L1

✓ PREV-12 Encounter, Outpatient or Office Visit

Encounter

Custom Codes

Encounter: OUTPATIENT, Encounter: OFFICE_VISIT

Procedure

CPT

HCPCS

SNOMED

90791, 90792, 90832, 90834, 90837, 90839, 92625, 96116, 96118, 96150, 96151, 97003, 99201, 99202, 99203, 99204, 99205, 99212, 99213, 99214, 99215, 99384, 99385, 99386, 99387, 99394, 99395, 99396, 99397	G0101, G0402, G0438, G0439, G0444	10197000, 108220007, 108221006, 108224003, 108250004, 108311000, 13607009, 14736009, 165171009, 171207006, 18512000, 185349003, 185463005, 185465003, 252592009, 252603000, 270427003, 270430005, 274803000, 277404009, 302440009, 30346009, 308335008, 32537008, 35025007, 370803007, 37894004, 390906007, 406547006, 410155007, 410157004, 43362002, 46662001, 48423005, 53555003, 67533008, 78318003, 83607001, 8411005, 86013001, 90526000, 91573000
---	-----------------------------------	--

Denominator Reference Tables

D1

Encounter

Numerator Reference Tables

N1	✓ depression screening				
Diagnosis					
ICD-9	ICD-10	SNOMED			
V79.0	Z13.89	428181000124104			
Observation					
LOINC					
73831-0, 73832-8					
Procedure					
HCPCS	SNOMED				
S3005	171207006				

Exclusions Reference Tables

XCL1	✓ Deceased		
	Death		
	Custom Codes		
	Deceased		
	Procedure		
	SNOMED		
	371828006		
XCL2	✓ Deceased		
	Admission		
XCL3	✓ Hospice		
	Encounter		
	Custom Codes		
	Encounter: type_hospice		
	Procedure		
	SNOMED		
	305336008, 385763009		
XCL4	✓ Dx of Depression or Bipolar Disorder in the Problem List		
	Diagnosis		
	ICD-9	ICD-10	SNOMED
	290.13, 290.21, 290.43, 296.20, 296.21, 296.22, 296.23, 296.24, 296.25, 296.26, 296.30, 296.31, 296.32, 296.33, 296.34, 296.36, 296.40, 296.41, 296.42, 296.43, 296.44, 296.45, 296.46, 296.80, 296.82, 298.0, 300.4, 301.12, 309.0, 309.1, 309.28, 311	F01.51, F31.11, F31.12, F31.13, F31.2, F31.73, F31.74, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.8, F32.9, F33.0, F33.1, F33.2, F33.3, F33.42, F33.9, F34.1, F43.21, F43.23	10875004, 10981006, 111485001, 1196001, 13313007, 13581000, 13746004, 14183003, 14495005, 1499003, 15193003, 15639000, 162004, 16295005, 16506000, 17782008, 18818009, 191610000, 191611001, 191613003, 191616006, 191618007, 191620005, 191621009, 191623007, 191625000, 191627008, 191629006, 191630001, 191634005, 191636007, 191638008, 191639000, 191641004, 191643001, 191659001, 192080009, 192362008, 19300006, 19527009, 25922000, 2618002, 26203008, 26530004, 268621008, 28475009, 28663008, 28884001, 29929003, 30520009, 30687003, 30935000, 3109008, 31446002, 319768000, 320751009, 33078009, 33380008, 34315001, 3530005, 35481005, 35489007, 35846004, 36170009, 36474008, 36583000, 36923009, 370143000, 371596008, 371599001, 371600003, 371604007, 38368003, 38451003, 38694004, 39809009, 40379007, 40568001, 40926005, 41552001, 41832009, 41836007, 42925002, 430852001, 43568002, 43769008, 442057004, 4441000, 45479006, 46229002, 48589009, 48937005, 49468007, 49512000, 51637008, 53049002, 53607008, 54761006, 55516002, 5703000, 59617007, 61403008, 63249007, 63778009, 64731001, 65042007, 66344007, 66631006, 67711008, 68569003, 69392006, 70546001, 71294008, 71336009, 71984005, 73471000, 73867007, 74686005, 75084000, 75360000, 75752004, 75837004, 76441001, 77486005, 77911002, 78269000, 78640000, 78667006, 79298009, 79584002, 81319007, 82998009, 83176005, 832007, 83225003, 84760002, 85080004, 85248005, 86058007, 87203005, 87512008, 87950005, 9340000

Exceptions Reference Tables

XCE1	✓ Medical or Patient Exception
	Diagnosis
	SNOMED
	105480006, 183932001, 183944003, 183945002, 397745006, 407563006, 413310006, 413311005, 413312003, 428119001

Reference Periods

START	STOP
12:00:01 AM	12:00:00 AM
01/01/2015	02/01/2015

Different criteria in each measure require that records occur during a precise time period. Note that all dates begin at midnight.

Operators

Operators describe what criteria is needed for a measure. They include:

AND

Both criteria above and below the line must be met

OR

Either criteria above or below the line must be met

NONE OF THE FOLLOWING

The criteria following this are negated — they must not have occurred

Expressions

You will encounter certain expressions that denote a specific relationship between criteria. They include:



Linked Encounter

Indicates that the criteria listed above must have occurred during the same encounter as the criteria that immediately follows

AND THEN

 HAPPENS AFTER

Indicates that the criteria must have occurred in a certain sequence. The criteria that follows must have occurred after the criteria listed above

AND THEN

 HAPPENED PREVIOUSLY

Reverse sequence — the criteria that follows must have occurred before the criteria listed above

PLUS

 COUNT INSTANCES

The first criteria specifies what must be on the patient record. The criteria below is then counted for that patient